arXiv:1204.5985v1 [math.PR] 26 Apr 2012

The Positive Occupation Time of Brownian Motion with Two-Valued Drift.

D.J.W. Simpson*
Department of Mathematics
University of British Columbia
Vancouver, BC
Canada

March 3, 2013

Abstract

The probability density function of the positive occupation time of one-dimensional Brownian motion with two-valued drift is derived by using the Feynman-Kac formula and Laplace transforms. The result is applied to a simple, two-dimensional stochastic differential equation describing stochastically perturbed sliding motion of a discontinuous, piecewise-smooth vector field. Long time asymptotics of the density are also computed.

1 Introduction

The one-dimensional stochastic differential equation

$$dx = \left\{ \begin{array}{ll} a_L \;, & x < 0 \\ -a_R \;, & x > 0 \end{array} \right\} dt + dW(t) \;, \qquad x(0) = x_0 \;, \tag{1}$$

where $a_L, a_R \in \mathbb{R}$ and $W(t)$ is a standard Brownian motion, represents *Brownian motion with two-valued drift*. Equation (1) has arisen in a stochastic control problem [1], investigations into the effects of noise on sliding motion in piecewise-smooth systems [2], and general theoretical settings [3, 4]. In the context of sliding motion it is assumed $a_L, a_R > 0$, such that solutions to (1) rarely stray far from the origin, but in this paper such a restriction is not made. An explicit expression for the PDF (probability density function) of $x(t)$ was first derived by Karatzas and Shreve in [5].

For any $t > 0$, let

$$\tau = \int_0^t \chi_{[0,\infty)}(x(s))\, ds = \mathrm{meas}\left\{ s \in [0,t] \;\middle|\; x(s) \geq 0 \right\} \;, \tag{2}$$

---

[*] The author thanks Rachel Kuske for many helpful discussions regarding this work.

denote the *positive occupation time* of $x(t)$. The purpose of this paper is to derive the PDF of τ, call it p, i.e.,

$$\mathrm{Prob}(\tau \in \Omega) = \int_{\Omega} p(\hat{\tau}; t; x_0, a_L, a_R)\, d\hat{\tau} \;, \tag{3}$$

for any measurable subset, $\Omega \subset [0, t]$. PDFs of occupation times of simple stochastic processes have found applications in control problems and mathematical finance [6].

The theorem below gives p for $x_0 = 0$. The result for $x_0 \neq 0$ may be written in terms of the result for $x_0 = 0$ by conditioning over the first passage time of $x(t)$ to zero. Specifically, let

$$h(s, x_0, a) = \frac{|x_0|}{\sqrt{2\pi s^3}}\, \mathrm{e}^{\frac{-(x_0 + as)^2}{2s}} \;, \tag{4}$$

denote the PDF for the first passage time, s, of $dx = a\, dt + dW(t)$, from $x(0) = x_0$ to $x(t) = 0$. Then

$$p(\tau; t; x_0, a_L, a_R) = \begin{cases} \int_0^t h(s; x_0, a_L) p(\tau; t-s; 0, a_L, a_R)\, ds + \delta(\tau) \int_t^\infty h(s, x_0, a_L)\, ds \;, & x_0 < 0 \\ \int_0^t h(s; x_0, -a_R) p(\tau - s; t-s; 0, a_L, a_R)\, ds + \delta(\tau - t) \int_t^\infty h(s, x_0, -a_R)\, ds \;, & x_0 > 0 \end{cases} \;, \tag{5}$$

where δ is the Dirac-delta function and we set $p(\tau; t; 0, a_L, a_R) = 0$ whenever $\tau \notin [0, t]$. For both $x_0 < 0$ and $x_0 > 0$, the first term of (5) corresponds to the case that $x(t)$ first reaches zero at a time $s \leq t$, and the second term corresponds to $x(t)$ not reaching zero by the time t.

Theorem 1.
For $x_0 = 0$ and any $t > 0$, the PDF of the positive occupation time (2) of Brownian motion with two-valued drift (1), is

$$\begin{aligned} p(\tau; t; 0, a_L, a_R) \;\; = \;\; & \frac{\mathrm{e}^{\frac{-a_L^2 (t-\tau)}{2}} \mathrm{e}^{\frac{-a_R^2 \tau}{2}}}{\pi \sqrt{\tau(t-\tau)}} - \frac{a_R\, \mathrm{e}^{\frac{-a_L^2 (t-\tau)}{2}} \operatorname{erfc}\left(\frac{a_R \sqrt{\tau}}{\sqrt{2}}\right)}{\sqrt{2\pi(t-\tau)}} - \frac{a_L\, \mathrm{e}^{\frac{-a_R^2 \tau}{2}} \operatorname{erfc}\left(\frac{a_L \sqrt{t-\tau}}{\sqrt{2}}\right)}{\sqrt{2\pi\tau}} \\ & + \frac{\sqrt{2}(a_L + a_R)}{\sqrt{\pi t}}\, \mathrm{e}^{\frac{-((a_L + a_R)\tau - a_L t)^2}{2t}} \operatorname{erfc}\left(\frac{-(a_L + a_R)\sqrt{\tau(t-\tau)}}{\sqrt{2t}}\right) \\ & + \mathcal{F}(\tau; t; a_L, a_R) + \mathcal{F}(t-\tau; t; a_R, a_L) \;, \end{aligned} \tag{6}$$

where

$$\mathcal{F}(\tau; t; a_L, a_R) = \frac{a_L(2a_L + a_R)}{2\sqrt{\pi}} \int_0^{t-\tau} -\frac{\sqrt{\tau}\mathrm{e}^{\frac{-a_R^2 \tau}{2}} \mathrm{e}^{\frac{-a_L^2 z}{2}}}{\sqrt{\pi z}(z+\tau)} + \frac{a_L z - a_R \tau}{\sqrt{2}(z+\tau)^{\frac{3}{2}}}\, \mathrm{e}^{\frac{-(a_L z - a_R \tau)^2}{2(z+\tau)}} \operatorname{erfc}\left(\frac{-(a_L + a_R)\sqrt{z\tau}}{\sqrt{2(z+\tau)}}\right) dz \;. \tag{7}$$

In two special cases, (6) reduces to known results. First, if $a_L = a_R = 0$, then $x(t)$ is simply regular Brownian motion. As first shown by Levy [7], in this case the occupation time is distributed by the arc-sine distribution [8, 9]. Indeed, in this case only the first term of (6) is nonzero, and

$$p(\tau; t; 0, 0, 0) = \frac{1}{\pi \sqrt{\tau(t-\tau)}} \;. \tag{8}$$

Equation (8) provides a reasonable approximation to $p(\tau;t;0,a_L,a_R)$ for small t, see Fig. 1, because over short time frames the diffusion of a stochastic quantity dominates its drift.

Secondly, if $a_L = -a_R$, then $x(t)$ is Brownian motion with constant drift. In this case (6) simplifies to

$$p(\tau;t;0,-a,a) = \left(\frac{1}{\sqrt{\pi\tau}} \mathrm{e}^{\frac{-a^2\tau}{2}} - \frac{a}{\sqrt{2}} \operatorname{erfc}\left(\frac{a\sqrt{\tau}}{\sqrt{2}} \right) \right) \left(\frac{1}{\sqrt{\pi(t-\tau)}} \mathrm{e}^{\frac{-a^2(t-\tau)}{2}} + \frac{a}{\sqrt{2}} \operatorname{erfc}\left(\frac{-a\sqrt{t-\tau}}{\sqrt{2}} \right) \right) , \tag{9}$$

which was first derived by Akatori [10]. The simplification is not straight-forward but may be demonstrated by using different expressions for Owen's T-function [11].

The remainder of this paper is organized as follows. A proof of Theorem 1 is presented in §2. The long time asymptotics of (6) are described in §3, focusing on the case $a_L, a_R > 0$. In §4, the result is applied to a model of stochastically perturbed sliding motion. Finally §5 contains a brief discussion.



Figure 1: The PDF (6) when $a_L = 2$ and $a_R = 1$. Note that the axes are scaled by t such that the domain is the unit interval at all times. For small t, p is well-approximated by the density of the arc-sine distribution, (8). For large t, p is approximately Gaussian since $a_L, a_R > 0$, see §3. When $t = 10$, on the given axes the Gaussian approximation has mean $\frac{a_L}{a_L+a_R} = \frac{2}{3}$ and standard deviation $\frac{1}{(a_L+a_R)\sqrt{t}} = \frac{1}{3\sqrt{10}} \approx 0.11$, as shown.

2 Proof of Theorem 1

For all $t > 0$, the stochastic differential equation (1) has a unique strong solution [12, 13] and therefore the expectation

$$u(x_0, t, \lambda) = \mathbb{E}_{x_0}\left(\mathrm{e}^{-\lambda \int_0^t \chi_{[0,\infty)}(x(s))\,ds}\right) , \tag{10}$$

where $\lambda > 0$, is well-defined. Equation (6) may be obtained from u because we have

$$u(0, t, \lambda) = \int_0^t \mathrm{e}^{-\lambda\tau} p(\tau; t; 0, a_L, a_R)\,d\tau \,. \tag{11}$$

Following a method applied to similar problems, see for instance [5, 10, 14], by the Feynman-Kac formula [8, 15], u is the unique, bounded, continuous solution to the initial value problem

$$u_t = \begin{cases} a_L u_{x_0} + \frac{1}{2} u_{x_0 x_0} \,, & x_0 < 0 \\ -a_R u_{x_0} + \frac{1}{2} u_{x_0 x_0} - \lambda u \,, & x_0 > 0 \end{cases} \,, \qquad u(x_0, 0, \lambda) = 1 \,. \tag{12}$$

Integration of the PDE with respect to x_0 over an arbitrarily small neighbourhood of zero reveals that u_{x_0} is also continuous.

To solve (12), we take the Laplace transform

$$U(x_0, \alpha, \lambda) = \int_0^\infty \mathrm{e}^{-\alpha t} u(x_0, t, \lambda)\,dt \,, \tag{13}$$

to produce the piecewise-linear ordinary differential equation

$$-1 = \begin{cases} \frac{1}{2} U_{x_0 x_0} + a_L U_{x_0} - \alpha U \,, & x_0 < 0 \\ \frac{1}{2} U_{x_0 x_0} - a_R U_{x_0} - (\alpha + \lambda) U \,, & x_0 > 0 \end{cases} \,. \tag{14}$$

Via standard ODE methods, we can obtain an explicit expression for the unique bounded solution to (14) for which u and u_{x_0} are continuous at $x_0 = 0$, and from this arrive at

$$U(0, \alpha, \lambda) = \frac{\alpha\left(\sqrt{2(\alpha+\lambda) + a_R^2} - a_R\right) + (\alpha+\lambda)\left(\sqrt{2\alpha + a_L^2} - a_L\right)}{\alpha(\alpha+\lambda)\left(\sqrt{2(\alpha+\lambda) + a_R^2} - a_R + \sqrt{2\alpha + a_L^2} - a_L\right)} \,. \tag{15}$$

Our goal is to obtain p, which by (11) and (13) is related to $U(0, \alpha, \lambda)$ by

$$U(0, \alpha, \lambda) = \int_0^\infty \int_0^t \mathrm{e}^{-\alpha t} \mathrm{e}^{-\lambda\tau} p(\tau; t; 0, a_L, a_R)\,d\tau\,dt \,. \tag{16}$$

Equation (16) is now used to derive p *constructively.* To make the task simply the evaluation of two inverse Laplace transforms, we first reverse the order of integration in (16), and let $v = t - \tau$ and $\mu = \alpha + \lambda$, to obtain

$$U(0, \alpha, \mu - \alpha) = \int_0^\infty \int_0^\infty \mathrm{e}^{-\alpha v} \mathrm{e}^{-\mu\tau} p(\tau, v + \tau; 0, a_L, a_R)\,dv\,d\tau \,. \tag{17}$$

Simplification is provided by noting that

$$U(0,\alpha,\mu-\alpha) = \Xi(\mu,\alpha;a_L,a_R) + \Xi(\alpha,\mu;a_R,a_L)\ , \tag{18}$$

where

$$\Xi(\mu,\alpha;a_L,a_R) = \frac{\sqrt{2\mu+a_R^2}-a_R}{\mu\left(\sqrt{2\mu+a_R^2}-a_R+\sqrt{2\alpha+a_L^2}-a_L\right)}\ . \tag{19}$$

By (17), with a little care it follows that

$$p(\tau;t;0,a_L,a_R) = \psi(\tau,t-\tau;a_L,a_R) + \psi(t-\tau,\tau;a_R,a_L)\ , \tag{20}$$

where

$$\Xi(\mu,\alpha;a_L,a_R) = \int_0^\infty\int_0^\infty \mathrm{e}^{-\alpha v}\mathrm{e}^{-\mu\tau}\psi(\tau,v;a_L,a_R)\,dv\,d\tau\ . \tag{21}$$

We now use (21) to obtain ψ from (19). Using, in particular, $\mathcal{L}^{-1}\left(\frac{1}{\sqrt{s}+k}\right) = \frac{1}{\sqrt{\pi t}} - k\mathrm{e}^{k^2 t}\,\mathrm{erfc}\left(k\sqrt{t}\right)$, the inverse Laplace transform of (19) with respect to α is

$$\begin{aligned}\int_0^\infty \mathrm{e}^{-\mu\tau}\psi(\tau,v;a_L,a_R)\,d\tau \quad &= \quad \frac{\sqrt{2\mu+a_R^2}-a_R}{2\mu}\,\mathrm{e}^{\frac{-a_L^2 v}{2}}\Bigg(\frac{\sqrt{2}}{\sqrt{\pi v}} - \left(\sqrt{2\mu+a_R^2}-a_R-a_L\right) \\ &\quad \times\ \mathrm{e}^{\left(\sqrt{2\mu+a_R^2}-a_R-a_L\right)^2\frac{v}{2}}\,\mathrm{erfc}\left(\left(\sqrt{2\mu+a_R^2}-a_R-a_L\right)\frac{\sqrt{v}}{\sqrt{2}}\right)\Bigg)\end{aligned} \tag{22}$$

To perform the daunting inverse Laplace transform of (22), we rewrite it as

$$\begin{aligned}\int_0^\infty \mathrm{e}^{-\mu\tau}\psi(\tau,v;a_L,a_R)\,d\tau \quad &= \quad \mathrm{e}^{\frac{-a_L^2 v}{2}}\Bigg(\frac{1}{\sqrt{\pi v}}\,G_1(\mu;a_R) - G_2(\mu,v;a_L,a_R) \\ &\quad + \frac{a_L+2a_R}{\sqrt{2}}\,G_1(\mu;a_R)G_2(\mu,v;a_L,a_R)\Bigg)\ ,\end{aligned} \tag{23}$$

where

$$G_1(\mu;a_R) \quad = \quad \frac{\sqrt{2\mu+a_R^2}-a_R}{\sqrt{2}\mu}\ , \tag{24}$$

$$G_2(\mu,v;a_L,a_R) \quad = \quad \mathrm{e}^{\left(\sqrt{2\mu+a_R^2}-a_R-a_L\right)^2\frac{v}{2}}\,\mathrm{erfc}\left(\left(\sqrt{2\mu+a_R^2}-a_R-a_L\right)\frac{\sqrt{v}}{\sqrt{2}}\right)\ , \tag{25}$$

so that

$$\psi(\tau,v;a_L,a_R) = \mathrm{e}^{\frac{-a_L^2 v}{2}}\left(\frac{1}{\sqrt{\pi v}}\,g_1(\tau;a_R) - g_2(\tau,v;a_L,a_R) + \frac{a_L+2a_R}{\sqrt{2}}\,g_1(\tau;a_R)*g_2(\tau,v;a_L,a_R)\right)\ , \tag{26}$$

where g_1 and g_2 denote the inverse Laplace transforms of G_1 and G_2 respectively, and $*$ denotes convolution with respect to τ. We have

$$g_1(\tau;a_R) = \frac{\mathrm{e}^{\frac{-a_R^2\tau}{2}}}{\sqrt{\pi\tau}} - \frac{a_R}{\sqrt{2}}\,\mathrm{erfc}\left(\frac{a_R\sqrt{\tau}}{\sqrt{2}}\right)\ , \tag{27}$$

and by the identity $\mathcal{L}^{-1}\left(\mathrm{e}^{(\sqrt{s}+k)^2}\operatorname{erfc}\left(\sqrt{s}+k\right)\right) = \frac{1}{2\pi t^{\frac{3}{2}}}\int_0^\infty z\mathrm{e}^{\frac{-z^2}{4}\left(1+\frac{1}{t}\right)}\mathrm{e}^{-kz}\,dz$ (obtained by combining formulas in [16]), we also have

$$g_2(\tau, v; a_L, a_R) = \frac{\mathrm{e}^{\frac{-a_R^2\tau}{2}}}{\sqrt{\pi}(\tau+v)}\left(\frac{\sqrt{v}}{\sqrt{\pi\tau}} + \frac{(a_L+a_R)v}{\sqrt{2(\tau+v)}}\mathrm{e}^{\frac{(a_L+a_R)^2\tau v}{2(\tau+v)}}\operatorname{erfc}\left(\frac{-(a_L+a_R)\sqrt{\tau v}}{\sqrt{2(\tau+v)}}\right)\right) . \quad (28)$$

After an involved algebraic simplification omitted for brevity, the combination of (20), (26), (27) and (28) yields (6) with (7) as required. ☐

3 Long time asymptotics of p

This section provides a statement of the long time asymptotics of (6), then a discussion of its derivation.

For large t,

$$p(\tau; t; 0, a_L, a_R) \sim \begin{cases} \frac{a_L+a_R}{\sqrt{2\pi t}}\,\mathrm{e}^{\frac{-((a_L+a_R)\tau - a_L t)^2}{2t}} , & a_L > 0,\ a_R > 0 \\ \mathcal{G}(\tau; a_L, a_R) , & a_L < 0,\ a_R > 0 \\ \mathcal{G}(t-\tau; a_R, a_L) , & a_L > 0,\ a_R < 0 \\ \mathcal{G}(\tau; a_L, a_R) + \mathcal{G}(t-\tau; a_R, a_L) , & a_L < 0,\ a_R < 0 \end{cases} , \quad (29)$$

where,

$$\mathcal{G}(\tau; a_L, a_R) = -\frac{\sqrt{2}a_L}{\sqrt{\pi\tau}}\,\mathrm{e}^{\frac{-a_R^2\tau}{2}} - a_L(2a_L+a_R)\,\mathrm{e}^{2a_L(a_L+a_R)\tau}\operatorname{erfc}\left(\frac{-(2a_L+a_R)\sqrt{\tau}}{\sqrt{2}}\right) . \quad (30)$$

To obtain (29), first note that it may be shown directly from (6), that for any fixed $\tau > 0$,

$$p(\tau; t; 0, a_L, a_R) \to \begin{cases} \mathcal{G}(\tau; a_L, a_R) , & a_L < 0 \\ 0 , & a_L > 0 \end{cases} , \quad (31)$$

as $t \to \infty$. For brevity we omit a derivation of this statement which requires taking $t \to \infty$ in (7) and simplifying the result via integral transformations. Moreover, for $a_L < 0$,

$$\int_0^\infty \mathcal{G}(\tau; a_L, a_R)\,d\tau = \begin{cases} \frac{a_L}{a_L+a_R} , & a_R \le 0 \\ 1 , & a_R \ge 0 \end{cases} , \quad (32)$$

which ensures (29) has unit area in the limit $t \to \infty$. The term $\mathcal{G}(t-\tau; a_R, a_L)$ appears in (29) by symmetry.

When $a_L, a_R > 0$, for large t the first three terms of (6) may be neglected because their L_1 norms rapidly approach zero as $t \to \infty$. The fourth term of (6) approaches $\frac{4(a_L+a_R)}{\sqrt{2\pi t}}\mathrm{e}^{\frac{-((a_L+a_R)\tau - a_L t)^2}{2t}}$, because here $\operatorname{erfc}(\cdot) \to 2$. To simplify (7), we consider the two terms in the integrand of (7) separately. The first term is negligible for large t when $a_L, a_R > 0$. By letting

$$\rho = t^\beta\left(\tau - \frac{a_L t}{a_L+a_R}\right) , \quad (33)$$

be a scaled difference of τ from the mean, $\frac{a_L t}{a_L+a_R}$, where $\beta \geq -1$ since $\tau \in [0,t]$, and substituting $u = \frac{a_L z - a_R \tau}{t}$, we have

$$\mathcal{F}(\tau;t;a_L,a_R) \to \frac{a_L^{\frac{3}{2}}(2a_L+a_R)\sqrt{t}}{2\sqrt{2\pi}} \int_{-\frac{a_L a_R}{a_L+a_R}-\frac{a_R\rho}{t^{\beta+1}}}^{-\frac{(a_L+a_R)\rho}{t^{\beta+1}}} \frac{u}{\left(a_L+\frac{(a_L+a_R)\rho}{t^{\beta+1}}+u\right)^{\frac{3}{2}}} \mathrm{e}^{\frac{-a_L t u^2}{2\left(a_L+\frac{(a_L+a_R)\rho}{t^{\beta+1}}+u\right)}} \operatorname{erfc}(\cdot)\, du \,. \tag{34}$$

Equation (34) may be evaluated asymptotically [17]. When $\rho \leq 0$, this is achieved by expanding the integrand in a Taylor series about $u = 0$; when $\rho > 0$, we expand about the upper limit of the integral. For instance, when $\rho \leq 0$ (the other case is similar and produces the same result), after integration we have

$$\mathcal{F}(\tau;t;a_L,a_R) \to -\frac{\sqrt{a_L}(2a_L+a_R)}{\sqrt{2\pi\left(a_L+\frac{(a_L+a_R)\rho}{t^{\beta+1}}\right)t}} \mathrm{e}^{\frac{-a_L(a_L+a_R)^2\rho^2}{2t^{2\beta+1}\left(a_L+\frac{(a_L+a_R)\rho}{t^{\beta+1}}\right)}} + O\left(\frac{1}{t}\right) \,. \tag{35}$$

Only when $\beta = -\frac{1}{2}$ is the leading order term of (35) non-constant in the limit $t \to \infty$, and in this case, after also substituting back τ using (33),

$$\mathcal{F}(\tau;t;a_L,a_R) \to -\frac{2a_L+a_R}{\sqrt{2\pi t}} \mathrm{e}^{\frac{-((a_L+a_R)\tau-a_L t)^2}{2t}} + O\left(\frac{1}{t}\right) \,. \tag{36}$$

Finally summing the non-neglected terms yields (29) for $a_L, a_R > 0$.

When $a_L, a_R > 0$, (29) is the PDF of a Gaussian with mean $\frac{a_L t}{a_L+a_R}$, and standard deviation $\frac{\sqrt{t}}{a_L+a_R}$, see Fig. 1. In this case the direction of the drift is towards the origin for both positive and negative x. The fraction of time spent in $[0,\infty)$, approaches $\frac{a_L}{a_L+a_R}$ as $t \to \infty$. This is consistent with the fact that the area under the steady-state density of $x(t)$ over $[0,\infty)$ is $\frac{a_L}{a_L+a_R}$, [2].

4 An application to stochastically perturbed sliding motion

Filippov systems are discontinuous vector fields used to model physical systems involving a discontinuity or abrupt change, such as mechanical systems with impacts and relay control systems [18, 19]. Stable *sliding motion* refers to dynamics that are constrained to a discontinuity surface due to a vector field that points towards the surface from both sides.

Consider the following two-dimensional stochastic differential equation,

$$\begin{bmatrix} dx_1 \\ dx_2 \end{bmatrix} = \left\{ \begin{array}{ll} \begin{bmatrix} a_L \\ b_L \end{bmatrix}, & x_1 < 0 \\ \begin{bmatrix} -a_R \\ b_R \end{bmatrix}, & x_1 > 0 \end{array} \right\} dt + \sqrt{\varepsilon} \begin{bmatrix} dW_1(t) \\ \sqrt{\kappa}\, dW_2(t) \end{bmatrix} \,, \tag{37}$$

where $W_1(t)$ and $W_2(t)$ are independent standard Brownian motions, $a_L, a_R > 0$, $b_L, b_R \in \mathbb{R}$ and $\varepsilon, \kappa \geq 0$. When $\varepsilon = 0$, (37) is Filippov and trajectories slide along $x_1 = 0$ because $a_L, a_R > 0$.

For small $\varepsilon > 0$, (37) describes stochastically perturbed sliding motion and $x_2(t)$ represents displacement along the discontinuity surface. We can write

$$x_2(t) = x_2(0) + b_L t + (b_R - b_L) \int_0^t \chi_{[0,\infty)}(x_1(s))\, ds + \sqrt{\varepsilon\kappa}\, W_2(t) \; . \tag{38}$$

Consequently the PDF of $x_2(t)$ is a convolution involving the PDF of the positive occupation time of $x_1(t)$ with the PDF of $W_2(t)$. Specifically, by using Theorem 1 and appropriately accommodating for ε, we have

$$\mathrm{Prob}(x_2(t) \in \Sigma) = \int_\Sigma \frac{1}{\varepsilon(b_R - b_L)} \int_{-\infty}^{\infty} p\left(\frac{\hat{x}_2 - x_2(0) - b_L t - w}{\varepsilon(b_R - b_L)} ; \frac{t}{\varepsilon} ; \frac{x_1(0)}{\varepsilon}, a_L, a_R \right) \frac{1}{\sqrt{2\pi\varepsilon\kappa t}}\, \mathrm{e}^{\frac{-w^2}{2\varepsilon\kappa t}}\, dw\, d\hat{x}_2 \; . \tag{39}$$

By (38), and the asymptotic calculations of §3, as $t \to \infty$,

$$\begin{aligned} \mathbb{E}(x_2(t)) &\to x_2(0) + b_L t + (b_R - b_L) \frac{a_L t}{a_L + a_R} = x_2(0) + \frac{a_L b_R + a_R b_L}{a_L + a_R} t \; , & (40) \\ \mathrm{Var}(x_2(t)) &\to \left(\frac{(b_R - b_L)^2}{(a_L + a_R)^2} + \kappa \right) \varepsilon t \; , & (41) \end{aligned}$$

which agrees with results of [2] that were obtained from the PDF of (1).

5 Discussion

This paper gives for the first time the PDF of the positive occupation time of Brownian motion with two-valued drift. The PDF, p, is given by the complicated expression (6). As $t \to \infty$, p approaches the simpler form (29). In the case that the piecewise drift points towards the origin on both sides, (29) is Gaussian. The results help us understand the system (37) that describes stochastically perturbed sliding motion of a piecewise-smooth vector field. An outstanding problem is the computation of the two-dimensional joint PDF of $x(t)$ and τ which would enable us to obtain the two-dimensional transitional PDF of (37).

References

[1] V.E. Beneš, L.A. Shepp, and H.S. Witsenhausen. Some solvable stochastic control problems. *Stochastics*, 4:39–83, 1980.

[2] D.J.W. Simpson and R. Kuske. Stochastically perturbed sliding motion in piecewise-smooth systems. Submitted to: *Proc. R. Soc. A*, 2012.

[3] M. Gradinaru, S. Herrmann, and B. Roynette. A singular large deviations phenomenon. *Ann. I. H. Poincaré*, 37(5):555–580, 2001.

[4] Z. Qian and W. Zheng. Sharp bounds for transition probability densities of a class of diffusions. *C.R. Acad. Sci. Paris, Ser. I*, 335:953–957, 2002.

[5] I. Karatzas and S.E. Shreve. Trivariate density of Brownian motion, its local and occupation times, with application to stochastic control. *Ann. Prob.*, 12(3):819–828, 1984.

[6] A. Pechtl. Distributions of occupation times of Brownian motion with drift. *J. Appl. Math. Decision Sci.*, 3(1):41–62, 1999.

[7] P. Lévy. Sur certains processus stochastiques homogènes. *Compositio Math.*, 7:283–339, 1939. In French.

[8] I. Karatzas and S.E. Shreve. *Brownian Motion and Stochastic Calculus.* Springer, New York, 1991.

[9] P. Billingsley. *Convergence of Probability Measures.* John Wiley & Sons, New York, 1999.

[10] J. Akahori. Some formulae for a new type of path-dependent option. *Ann. Appl. Probab.*, 5(2):383–388, 1995.

[11] D.B. Owen. Tables for computing bivariate normal probabilities. *Ann. Math. Stat.*, 27:1075–1090, 1956.

[12] D. Stroock and S.R.S Varadhan. Diffusion processes with continuous coefficients. I. *Comm. Pure Appl. Math.*, 22:345–400, 1969.

[13] Yu.V. Prokhorov and A.N. Shiryaev, editors. *Probability Theory III: Stochastic Calculus.* Springer, New York, 1998.

[14] J.M. Steele. *Stochastic Calculus and Financial Applications.* Springer, New York, 2001.

[15] B. Øksendal. *Stochastic Differential Equations: An Introduction with Applications.* Springer, New York, 2003.

[16] G.E. Roberts and H. Kaufman. *Table of Laplace Transforms.* W. B. Saunders Company, Philadelphia, 1966.

[17] C.M. Bender and S.A. Orszag. *Advanced Mathematical Methods for Scientists and Engineers.* International Series in Pure and Applied Mathematics. McGraw-Hill, New York, 1978.

[18] M. di Bernardo, C.J. Budd, A.R. Champneys, and P. Kowalczyk. *Piecewise-smooth Dynamical Systems. Theory and Applications.* Springer-Verlag, New York, 2008.

[19] A.F. Filippov. *Differential Equations with Discontinuous Righthand Sides.* Kluwer Academic Publishers., Norwell, 1988.